## CORRECTION Open Access



## Correction: Median nerve block increases the success rate of radial artery cannulation in women with gestational hypertension undergoing cesarean section

Xin Men<sup>1</sup>, Qian Wang<sup>2</sup>, Wen-sheng Hu<sup>3</sup>, Yun Chai<sup>3</sup>, Ting-ting Ni<sup>4</sup>, Hong-yan Shou<sup>1\*†</sup> and Zhen-feng Zhou<sup>1\*†</sup>

Correction: BMC Anesthesiol 22, 248 (2022) https://doi.org/10.1186/s12871-022-01793-4

Following publication of the original article [1], the authors identified an error in the author names of Wensheng Hu and Hong-yan Shou.

The incorrect author names are: Wen-shen Hu and Hong-ye Sho

The correct author names are: Wen-sheng Hu and Hong-yan Shou

The author group has been updated above and the original article [1] has been corrected.

## **Author details**

<sup>1</sup>Department of Anesthesiology, Hangzhou Women's Hospital (Hangzhou Maternity and Child Health Care Hospital, Hangzhou First People's Hospital Qianjiang New City Campus, The Affiliated Women's Hospital of Hangzhou Normal University), Hangzhou 315014, China. <sup>2</sup>Department of Anesthesiology, The Affiliated ZheJiang Hospital, School of Medicine, Zhejiang University, Hangzhou 315014, China. <sup>3</sup>Department of Obstetrics, Hangzhou Women's Hospital (Hangzhou Maternity and Child Health Care Hospital, Hangzhou First People's Hospital Qianjiang New City Campus, The Affiliated Women's Hospital

The original article can be found online at https://doi.org/10.1186/s12871-022-01793-4.

<sup>†</sup>Hong-yan Shou and Zhen-feng Zhou contributed equally to this work.\*Correspondence: shouhongyan66@163.com; zhenfeng9853@163.com

of Hangzhou Normal University), Hangzhou 315014, China. <sup>4</sup>Department of Anesthesiology, Ningbo NO.7 Hospital, Ningbo 320000, China.

Published online: 27 August 2022

## Reference

 Men X, Wang Q, Hu W, et al. Median nerve block increases the success rate of radial artery cannulation in women with gestational hypertension undergoing cesarean section. BMC Anesthesiol. 2022;22:248. https://doi. org/10.1186/s12871-022-01793-4.



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and given intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativeccommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

<sup>&</sup>lt;sup>1</sup> Department of Anesthesiology, Hangzhou Women's Hospital (Hangzhou Maternity and Child Health Care Hospital, Hangzhou First People's Hospital Qianjiang New City Campus, The Affiliated Women's Hospital of Hangzhou Normal University), Hangzhou 315014, China Full list of author information is available at the end of the article