

CORRECTION

Open Access



# Correction: Motor-sparing peripatellar plexus block provides noninferior block duration and complete block area of the peripatellar region compared with femoral nerve block: a randomized, controlled, noninferiority study

Wen-Yi Gong<sup>1</sup>, Chen-Guang Li<sup>2†</sup>, Jing-Yu Zhang<sup>3†</sup>, Xiao-Hui Liao<sup>4</sup>, Cheng Zhu<sup>4</sup>, Jie Min<sup>4</sup>, Xiao-Fang Yue<sup>5\*</sup> and Kun Fan<sup>6\*</sup>

**Correction: BMC Anesthesiol 22, 334 (2022)**  
<https://doi.org/10.1186/s12871-022-01863-7>

Published online: 09 December 2022

Following publication of the original article [1], the authors identified an error in Fig. 4. The correct figure is given below.

The original article [1] has been updated.

#### Author details

<sup>1</sup>Department of Anaesthesiology, Wusong Central Hospital, Shanghai, China. <sup>2</sup>Department of Anaesthesiology, First People's Hospital of Tianshui, Gansu, China. <sup>3</sup>Department of Anaesthesiology, Second Hospital Affiliated to Lanzhou University, Gansu, China. <sup>4</sup>Department of Orthopaedics, Wusong Central Hospital, Shanghai, China. <sup>5</sup>Department of Neurology, Shanghai Sixth People's Hospital, No. 600, Yishan Road, Shanghai 200233, China. <sup>6</sup>Department of Anaesthesiology, Shanghai Sixth People's Hospital, No. 600, Yishan Road, Shanghai 200233, China.

The original article can be found online at <https://doi.org/10.1186/s12871-022-01863-7>.

<sup>†</sup>Chen-Guang Li and Jing-Yu Zhang contributed equally to this work.\*Correspondence: [klyxf@126.com](mailto:klyxf@126.com); [fankunmazui@hotmail.com](mailto:fankunmazui@hotmail.com)

<sup>5</sup> Department of Neurology, Shanghai Sixth People's Hospital, No. 600, Yishan Road, Shanghai 200233, China

<sup>6</sup> Department of Anaesthesiology, Shanghai Sixth People's Hospital, No. 600, Yishan Road, Shanghai 200233, China


Full list of author information is available at the end of the article

#### Reference

1. Gong WY, Li CG, Zhang JY, et al. Motor-sparing peripatellar plexus block provides noninferior block duration and complete block area of the peripatellar region compared with femoral nerve block: a randomized, controlled, noninferiority study. *BMC Anesthesiol.* 2022;22:334. <https://doi.org/10.1186/s12871-022-01863-7>.



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.



**Fig. 4** Course and staining of the peripatellar plexus 30 min after dye injection. **A** Course and staining of the LFCN; **B** Course and staining of the IPBSN; **C** Course and staining of the IFCN; **D** Course and staining of the aMFCN; **E** Staining of the FN. ASIS, anterior superior iliac spine; SM, sartorius muscle; aLFCN, anterior branch of the lateral femoral cutaneous nerve; pLFCN, posterior branch of the lateral femoral cutaneous nerve; IPBSN, infrapatellar branch of the saphenous nerve; IFCN I, intermediate femoral cutaneous nerve branch I; IFCN II, intermediate femoral cutaneous nerve branch II; aMFCN, anterior branch of the medial femoral cutaneous nerve; SN, saphenous nerve; FN, femoral nerve