CORRECTION   **Open Access**

Correction: Rate of residual neuromuscular block using single-dose rocuronium in general anesthesia for ENT Surgery: a retrospective cohort study

Orlando Carrillo-Torres[1], María Guadalupe Pliego-Sánchez[1], Víctor Joshua Pérez-Muñoz[1], Jennifer Sánchez-Jurado[1], Verónica Camacho-Vacherón[1] and José Damián Carrillo-Ruiz[2,3]*

Correction: BMC Anesthesiol 23, 107 (2023).
https://doi.org/10.1186/s12871-023-02027-x.

Following publication of the original article [1], the authors reported that the last names of the authors are not correct as parts of the family names were captured as given names. The correct presentation of the authors names are as follows:

 Orlando Carrillo-Torres
María Guadalupe Pliego-Sánchez
Victor Joshua Pérez-Muñoz
Jennifer Sánchez-Jurado
Verónica Camacho-Vacherón
José Damián Carrillo-Ruíz

The author group has been updated above and the original article [1] has been corrected.

The online version of the original article can be found at https://doi.org/10.1186/s12871-023-02027-x.

*Correspondence:
José Damián Carrillo-Ruiz
josecarrilloruiz@yahoo.com
[1]Anesthesiology Service at Mexico General Hospital, Mexico City, Mexico
[2]Research Direction & Neurosurgery Service at Hospital General de México, Mexico City, Mexico
[3]Neuroscience Coordination of Psychology Faculty at Mexico Anahuac University, Av. Anahuac 46, Lomas Anahuac, Naucalpan de Juárez, Estado de México 52786, Mexico

Published online: 19 October 2023

References
1. Carrillo-Torres O, Pliego-Sánchez MG, Pérez-Muñoz VJ, et al. Rate of residual neuromuscular block using single-dose rocuronium in general anesthesia for ENT Surgery: a retrospective cohort study. BMC Anesthesiol. 2023;23:107. https://doi.org/10.1186/s12871-023-02027-x.

Publisher's Note
Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.