CORRECTION

Correction: Impact of total intravenous anesthesia and total inhalation anesthesia as the anesthesia maintenance approaches on blood glucose level and postoperative complications in patients with type 2 diabetes mellitus: a double-blind, randomized controlled trial

Xinghui Xiong^{1†}, Yong He^{2†}, Cheng Zhou³, Qin Zheng², Chan Chen^{1,3*} and Peng Liang^{1,2,4*}

Correction: BMC Anesthesiol 23, 267 (2023) https://doi.org/10.1186/s12871-023-02199-6

Following publication of the original article [1], the authors reported an error found under "Intervention and measurement" section. The dose of propofol in continuous infusion used in the TIVA group was $4-12 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ instead of $4-12 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$.

[†]Xinghui Xiong and Yong He contributed equally to this work.

The online version of the original article can be found at https://doi.org/10.1186/s12871-023-02199-6.

*Correspondence: Chan Chen xychenchan@gmail.com Peng Liang liangpengwch@scu.edu.cn ¹Department of Anesthesiology, West China Hospital, Sichuan University, Chengdu 610041, Sichuan, China ²Department of Laboratory Medicine, West China Hospital, Sichuan University, Chengdu 610041, Sichuan, China ³Laboratory of Anesthesia and Critical Care Medicine, West China Hospital, National-Local Joint Engineering Research Centre of Translational Medicine of Anesthesiology, Sichuan University, Chengdu 610041, Sichuan, China ⁴Day Surgery Center, West China Hospital, Sichuan University,

Chengdu 610041, Sichuan, China

© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

The original article [1] has been updated.

Published online: 22 January 2024

References

 Xiong X, He Y, Zhou C, et al. Impact of total intravenous anesthesia and total inhalation anesthesia as the anesthesia maintenance approaches on blood glucose level and postoperative complications in patients with type 2 diabetes mellitus: a double-blind, randomized controlled trial. BMC Anesthesiol. 2023;23:267. https://doi.org/10.1186/s12871-023-02199-6.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.







