

Research article

Open Access

## High frequency oscillatory ventilation and prone positioning in a porcine model of lavage-induced acute lung injury

Joerg Brederlau\*<sup>1</sup>, Ralf Muellenbach<sup>1</sup>, Markus Kredel<sup>1</sup>, Clemens Greim<sup>2</sup> and Norbert Roewer<sup>1</sup>

Address: <sup>1</sup>Klinik und Poliklinik für Anästhesiologie, Universitätsklinikum Würzburg, Oberdürrbacher Str. 6, 97080 Würzburg, Germany and <sup>2</sup>Klinik für Anästhesiologie, Intensiv- und Notfallmedizin, Klinikum Fulda, Pacelliallee 4, 36043 Fulda, Germany

Email: Joerg Brederlau\* - brederlau\_j@klinik.uni-wuerzburg.de; Ralf Muellenbach - muellenbach\_r@klinik.uni-wuerzburg.de; Markus Kredel - kredel\_m@klinik.uni-wuerzburg.de; Clemens Greim - greim@klinikum-fulda.de; Norbert Roewer - roewer\_n@klinik.uni-wuerzburg.de

\* Corresponding author

Published: 03 April 2006

Received: 22 November 2005

BMC Anesthesiology 2006, 6:4 doi:10.1186/1471-2253-6-4

Accepted: 03 April 2006

This article is available from: <http://www.biomedcentral.com/1471-2253/6/4>

© 2006 Brederlau et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** This animal study was conducted to assess the combined effects of high frequency oscillatory ventilation (HFOV) and prone positioning on pulmonary gas exchange and hemodynamics.

**Methods:** Saline lung lavage was performed in 14 healthy pigs ( $54 \pm 3.1$  kg, mean  $\pm$  SD) until the arterial oxygen partial pressure ( $\text{PaO}_2$ ) decreased to  $55 \pm 7$  mmHg. The animals were ventilated in the pressure controlled mode (PCV) with a positive endexpiratory pressure (PEEP) of 5 cmH<sub>2</sub>O and a tidal volume ( $V_T$ ) of 6 ml/kg body weight. After a stabilisation period of 60 minutes, the animals were randomly assigned to 2 groups. Group 1: HFOV in supine position; group 2: HFOV in prone position. After evaluation of prone positioning in group 2, the mean airway pressure ( $P_{\text{mean}}$ ) was increased by 3 cmH<sub>2</sub>O from 16 to 34 cmH<sub>2</sub>O every 20 minutes in both groups accompanied by measurements of respiratory and hemodynamic variables. Finally all animals were ventilated supine with PCV, PEEP = 5 cm H<sub>2</sub>O,  $V_T$  = 6 ml/kg.

**Results:** Combination of HFOV with prone positioning improves oxygenation and results in normalisation of cardiac output and considerable reduction of pulmonary shunt fraction at a significant ( $p < 0.05$ ) lower  $P_{\text{mean}}$  than HFOV and supine positioning.

**Conclusion:** If ventilator induced lung injury is ameliorated by a lower  $P_{\text{mean}}$ , a combined treatment approach using HFOV and prone positioning might result in further lung protection.

### Background

Prevention of irreversible hypoxemic damage and improvement of respiratory mechanics are the main treatment goals in patients with acute respiratory distress syndrome (ARDS). Mechanical ventilation is the predominant supportive treatment modality in ARDS, but

has also detrimental side effects, currently termed ventilator induced lung injury (VILI).

Lung protective ventilation strategies aim to ameliorate VILI by application of a reduced tidal volume ( $V_T$  = 6 ml/

kg), sufficient PEEP-level and limitation of the plateau inspiratory pressure to 35 cm H<sub>2</sub>O [1-3].

Although it is known, that the degree of hypoxemia is inconclusive to predict mortality [4], the early response of the PaO<sub>2</sub>/FIO<sub>2</sub>-ratio to therapeutic interventions might be an indicator for an increased survival rate in ARDS [5,6]. This calls for the most rapid amelioration of hypoxemia with a mono- or multimodal organ protective treatment approach.

High frequency oscillatory ventilation (HFOV) with its constant mean airway pressure ( $P_{\text{mean}}$ ) with superimposed small tidal volumes and active in- and expiration at a high respiratory frequency might be the ideal lung protective ventilatory strategy [7]. In a multicenter randomized controlled trial investigating the effectiveness of HFOV, the significant early improvement of the PaO<sub>2</sub>/FIO<sub>2</sub>-ratio in the HFOV-group was associated with a tendency towards a reduced 30-day mortality compared with the conventional ventilation group. The PaO<sub>2</sub>/FIO<sub>2</sub>-ratio was the most significant predictor of survival independent of the selected ventilator strategy [8].

Prone positioning was shown to increase the PaO<sub>2</sub> in 70–80% of patients with ARDS and to improve alveolar ventilation without influencing the 28-day mortality [9,10]. If PaCO<sub>2</sub>-reduction was achievable with prone positioning, 28-day mortality in ARDS patients was significantly reduced [11]. Prone positioning and application of PEEP were shown to have an additive effect on oxygenation [12]. However, prone positioning is a potentially dangerous manoeuvre with acute and long term complications such as tracheal tube dislocation, and pressure sores [13].

Combination of different treatments are used in desperation for salvage therapy in patients with ARDS [14]. A recently published study in 39 medical ARDS-patients randomized to conventional lung protective ventilation and HFOV showed comparable increases of the PaO<sub>2</sub>/FIO<sub>2</sub>-ratio after prone positioning. An additive effect of prone positioning and HFOV could not be demonstrated [15].

The objective of our study was to evaluate the effects of prone positioning on gas-exchange, hemodynamics and respiratory parameters in HFOV-ventilated pigs with severe lavage induced acute lung injury [16]. We hypothesized, that during HFOV oxygenation can be improved at a lower  $P_{\text{mean}}$  with the animals positioned prone than supine.

## Methods

### Animals

The study was conducted in accordance with the National Institutes of Health guidelines for ethical animal research and was approved by the Laboratory Animal Care and Use Committee of the District of Unterfranken, Germany.

The experiments were performed in 14 healthy pigs, Pietrain breed, all negative for the malignant hyperthermia gene. The animals were 14 to 18 weeks old, with a mean ( $\pm$  SD) body weight of 54  $\pm$  3.1 kg.


### Experimental preparation

The animals were fasted for 24 hours without limiting water access. Prior to instrumentation the animals were sedated with intramuscular ketamin (10 mg/kg), xylazine hydrochloride (1 mg/kg) and atropine (25  $\mu$ g/kg) and placed supine on an operating table armed with a heating pad to provide core temperature stability (37.3  $\pm$  0.5 °C). Anesthesia was induced with an intravenous bolus of sodium thiopental (5 mg/kg) using an auricular vein. The animals' trachea was orally intubated with a cuffed 8.0-mm ID Edgar tracheal tube (Rueschelit<sup>®</sup>, Ruesch, Kernen, Germany) providing an additional lumen embedded in the tubes inner wall for tracheal pressure monitoring. Anesthesia and complete muscle relaxation were maintained with continuous intravenous infusion of ketamin (2 mg/kg/h), midazolam (0.5 mg/kg/h), fentanyl (0.01 mg/kg/h) and vecuronium (0.1 mg/kg/h).

The animals were mechanically ventilated with a Servo<sup>®</sup> 900C ventilator (Siemens-Elema AB, Solna, Sweden) using pressure controlled ventilation (PCV) with a PEEP of 5 cmH<sub>2</sub>O, an inspiratory to expiratory ratio (I:E) of 1:1 and a fraction of inspired oxygen (FIO<sub>2</sub>) of 1.0. A V<sub>T</sub> of 6 ml/kg and a respiratory rate (RR) of 25–30 breath/min were applied resulting in normocapnia.

After a bolus of 500 ml balanced electrolyte solution a continuous infusion was given at a rate of 2–6 ml/kg/h. Continuous electrocardiography (Servomed<sup>®</sup>, Hellige, Freiburg i. Br., Germany), pulseoxymetry, capnography and distal tracheal pressure monitoring (SM8050<sup>®</sup>, Draeger, Luebeck, Germany) were performed.

2 gm Cefotiam was administered intravenously. After systemic heparinization (300 U/kg Liquemin<sup>®</sup>, Roche, Reinach, Switzerland) arterial and central venous access were established transcutaneously using ultrasound guidance (SonoSite 180 Plus<sup>®</sup>, SonoSite Inc., Botell, WA, USA). Activated clotting time (ACT II<sup>®</sup>, Medtronic, Minneapolis, MN, USA) was measured hourly and maintained between 150 and 200 seconds throughout the experiment with heparin bolus injections as needed. The left carotid artery was cannulated with a 20-gauge catheter (Vygon, Ecouen,



**Figure 1**

Experimental protocol  $T_{baseline}$ : 30 min after instrumentation.  $T_0$ : 60 min after last pulmonary lavage.  $T_{20}$ : HFOV ( $P_{mean} = 16$  cmH<sub>2</sub>O).  $T_{40}$ : prone positioning in group 2.  $T_{60}$ :  $P_{mean}$  increased from 16 to 19 cmH<sub>2</sub>O.  $T_{80}$ :  $P_{mean} = 22$  cmH<sub>2</sub>O.  $T_{100}$ :  $P_{mean} = 25$  cmH<sub>2</sub>O.  $T_{120}$ :  $P_{mean} = 28$  cmH<sub>2</sub>O.  $T_{140}$ :  $P_{mean} = 31$  cmH<sub>2</sub>O.  $T_{160}$ :  $P_{mean} = 34$  cmH<sub>2</sub>O.  $T_{180}$ :  $P_{mean} = 16$  cmH<sub>2</sub>O ( $\approx T_0$ ) PCV in all groups with HFOV and prone positioning discontinued.

France). The right internal jugular vein was cannulated with a 9 French introducer sheath (Arrow, Reading, PA, USA) and a 7,5 French flow directed thermodilution pulmonary artery catheter (831F75, Edwards Lifescience, Irvine, CA, USA) was advanced into the pulmonary artery under transduced pressure guidance.

**Hemodynamic, ventilatory and blood gas measurements**

For hemodynamic monitoring pressure transducers referenced to atmospheric pressure at the mid-thoracic level (Combitrans®, Braun, Melsungen, Germany) and a modular monitor system (Servomed®, Hellige, Freiburg i. Br., Germany) were used. Mean arterial pressure (MAP), mean pulmonary artery pressure (MPAP), central venous pressure (CVP) and pulmonary artery occlusion pressure (PCWP) were transduced. Heart rate (HR) was traced by the electrocardiogram.

Trifold injections of 10-ml aliquots of ice cold saline into the right atrium at random phases of different respiratory cycles were used for pulmonary artery catheter-based cardiac output (CO)-measurements (Explorer®, Edwards Lifescience, Irvine, CA, USA).

Arterial and mixed venous blood samples were immediately analyzed for PO<sub>2</sub>, PCO<sub>2</sub> and pH using standard blood gas electrodes (ABL 505®, Radiometer, Bronshoj,

Denmark). In each sample, hemoglobin and oxygen saturation were measured using spectrophotometry (OSM3®, Radiometer, Bronshoj, Denmark). Arterial (CaO<sub>2</sub>), mixed venous (CvO<sub>2</sub>) and pulmonary capillary (CCO<sub>2</sub>) oxygen contents (ml/dl) and the pulmonary shunt fraction (Qs/Qt) were calculated using standard formulas. The oxygenation index (OI) was calculated using the formula introduced by Hallmann et al.:  $OI = (P_{mean} \times FiO_2 \times 100) / PaO_2$  [17].

For tracheal pressure monitoring air filled pressure transducers (Combitrans®, Braun, Melsungen, Germany) referenced to atmospheric pressure were used [18]. Temperature was measured by thermistor in the pulmonary artery.

**Experimental procedure**

*Lung injury*

After instrumentation the animals were stabilized for 30 min in the supine position and mechanically ventilated with PCV ( $V_T = 6$  ml/kg, I:E = 1:1,  $FiO_2 = 1.0$ , PEEP = 5 cmH<sub>2</sub>O). RR was adjusted to achieve normocapnia. Baseline measurements were obtained.

Lung injury was induced by bilateral pulmonary lavages with 30 ml/kg isotonic saline (38°C) and repeated every 10 minutes until PaO<sub>2</sub> decreased to 40-60 mmHg and

**Table 1: Hemodynamic and metabolic data at baseline (T<sub>baseline</sub>), after injury (T<sub>0</sub>), after starting HFOV (T<sub>20</sub>), after randomisation (T<sub>40</sub>), during P<sub>mean</sub>-augmentation (T<sub>60</sub>-T<sub>160</sub>), at end of experiment (T<sub>180</sub>)**

	Group	T <sub>baseline</sub>	T <sub>0</sub>	T <sub>20</sub>	T <sub>40</sub>	T <sub>60</sub>	T <sub>80</sub>	T <sub>100</sub>	T <sub>120</sub>	T <sub>140</sub>	T <sub>160</sub>	T <sub>180</sub>
PaCO <sub>2</sub> [mmHg]	HFOV	41 +/- 3 #	76 +/- 12 *	73 +/- 8	73 +/- 8	74 +/- 9	74 +/- 10	78 +/- 12	80 +/- 12	82 +/- 12	83 +/- 13	85 +/- 10 #
	HFOV prone	41 +/- 3 #	77 +/- 6 *	79 +/- 6	70 +/- 6	71 +/- 6	72 +/- 6	75 +/- 4	76 +/- 4	77 +/- 5	77 +/- 6	78 +/- 5
pH	HFOV	7,43 +/- 0,05 #	7,11 +/- 0,13 *§	7,15 +/- 0,09	7,15 +/- 0,09	7,19 +/- 0,1	7,21 +/- 0,09 #	7,21 +/- 0,1 #	7,15 +/- 0,13	7,15 +/- 0,13	7,15 +/- 0,13	7,16 +/- 0,12
	HFOV prone	7,45 +/- 0,03 #	7,22 +/- 0,1 *	7,19 +/- 0,09	7,2 +/- 0,06	7,23 +/- 0,06	7,21 +/- 0,05	7,21 +/- 0,05	7,19 +/- 0,06	7,18 +/- 0,07	7,19 +/- 0,07	7,22 +/- 0,1
Qs/Qt (ratio)	HFOV	0,01 +/- 0,03 #	0,65 +/- 0,15 *	0,57 +/- 0,12	0,57 +/- 0,12 §	0,5 +/- 0,16 #§	0,42 +/- 0,21 *#§	0,33 +/- 0,26 *#§	0,24 +/- 0,2 *	0,19 +/- 0,15 #	0,15 +/- 0,1 #	0,64 +/- 0,09 *
	HFOV prone	0 +/- 0,02 #	0,56 +/- 0,07*	0,5 +/- 0,08	0,36 +/- 0,11 *#	0,24 +/- 0,13 *#	0,2 +/- 0,1 #	0,16 +/- 0,1 #	0,15 +/- 0,1 #	0,13 +/- 0,09 #	0,12 +/- 0,06 #	0,56 +/- 0,05 *
SgvO <sub>2</sub> [%]	HFOV	79 +/- 10 #	63 +/- 10 *§	57 +/- 8	57 +/- 8	61 +/- 11	65 +/- 11	69 +/- 7	72 +/- 3	75 +/- 2 #	76 +/- 3 #	53 +/- 11 #*
	HFOV prone	87 +/- 3 #	48 +/- 10 *	53 +/- 8	65 +/- 14 *#	67 +/- 5 #	73 +/- 4 #	71 +/- 6 #	72 +/- 7 #	74 +/- 3 #	75 +/- 4 #	47 +/- 11 *
PIP [cmH <sub>2</sub> O]	HFOV	20 +/- 3 #	28 +/- 3 *	20 +/- 1 *#	20 +/- 1 #	23 +/- 1 *#	26 +/- 1 *#	28 +/- 1 *	32 +/- 1 *#	35 +/- 1 *#	37 +/- 1 *#	27 +/- 3 *§
	HFOV prone	18 +/- 3 #	28 +/- 2 *	20 +/- 1 *#	20 +/- #	23 +/- 1 *#	26 +/- 1 *	28 +/- 1	32 +/- 1 *#	35 +/- 1 *#	37 +/- 1 *#	29 +/- 2 *
HR [1/min]	HFOV	87 +/- 19	89 +/- 14 §	79 +/- 15	80 +/- 15 §	78 +/- 18 §	75 +/- 18 §	78 +/- 21 §	75 +/- 19 §	69 +/- 12 #	76 +/- 21	77 +/- 18
	HFOV prone	83 +/- 15 #	66 +/- 13 *	68 +/- 15	58 +/- 8	57 +/- 8	53 +/- 4	52 +/- 5	56 +/- 4	57 +/- 4	62 +/- 9	75 +/- 11
MAP [mmHg]	HFOV	79 +/- 13	89 +/- 11	81 +/- 12	81 +/- 12	81 +/- 8	85 +/- 8	84 +/- 5	84 +/- 5	82 +/- 5	84 +/- 9	84 +/- 4
	HFOV prone	83 +/- 9	84 +/- 7	83 +/- 7	83 +/- 10	81 +/- 8	79 +/- 8	80 +/- 7	85 +/- 12	78 +/- 5	78 +/- 8	83 +/- 7
MPAP [mmHg]	HFOV	23 +/- 5 #	32 +/- 6 *	33 +/- 6	33 +/- 6	33 +/- 6	34 +/- 5	37 +/- 6	37 +/- 4 #	38 +/- 4 #	39 +/- 4 #	33 +/- 5
	HFOV prone	22 +/- 6 #	29 +/- 4 *	34 +/- 4	32 +/- 6	35 +/- 5 #	34 +/- 3	35 +/- 3 #	35 +/- 4 #	36 +/- 2 #	36 +/- 2 #	30 +/- 4 *
CVP [mmHg]	HFOV	6,7 +/- 1,3 #	9 +/- 2,4 *	9,7 +/- 2,5	9,7 +/- 2,5	10 +/- 2,8 §	11,6 +/- 2,9 #	12 +/- 2,4 #	13,7 +/- 1,3 #	13,9 +/- 0,9 #	14,3 +/- 1,1 #	9,4 +/- 1,3 *
	HFOV prone	6,9 +/- 1,3 #	10,9 +/- 3,3 *	10 +/- 1	10,9 +/- 1,5	12,7 +/- 2,9	11,9 +/- 2,3	12,4 +/- 2,4	12,3 +/- 2,6	12,6 +/- 2,1	12,6 +/- 2,1	10,7 +/- 2,3
PCWP [mmHg]	HFOV	9,1 +/- 2,3	10 +/- 3,7	11 +/- 3,8	11 +/- 3,8	12 +/- 3,2	12,1 +/- 2,6	12,4 +/- 2,2	14,1 +/- 2,4 #	15 +/- 1,9 #	15 +/- 1,8 #	11 +/- 3,3 *
	HFOV prone	9,1 +/- 2,3	11,1 +/- 1,9	11,9 +/- 1,3	11,3 +/- 2,1	12,9 +/- 2,7	12,6 +/- 1,7	12,9 +/- 1,9	12,1 +/- 1,6	12,9 +/- 1,2	12,6 +/- 1,1	11,1 +/- 1,3

Data are mean ± standard deviation; Two-way-ANOVA with repeated measurements (Student-Newman-Keuls Method – post hoc test): \* p < 0.05 vs. T<sub>baseline</sub>; # p < 0.05 vs. T<sub>0</sub>; § p < 0.05 HFOV vs. HFOV prone. CVP = central venous pressure; HFOV = high frequency oscillatory ventilation; HR = heart rate; MAP = mean arterial pressure; MPAP = mean pulmonary artery pressure; PCV = pressure controlled ventilation; PCWP = pulmonary capillary wedge pressure; Qs/Qt = pulmonary shunt fraction; SgvO<sub>2</sub> = mixed venous oxygen saturation.

was stable for 60 minutes with unchanged ventilator parameters. During induction of lung injury all lungs were ventilated with PCV, FIO<sub>2</sub> = 1.0, PEEP = 5 cmH<sub>2</sub>O, V<sub>T</sub> = 6 ml/kg, RR = 40/min. Post injury measurements were obtained.

**Positioning**

Prone positioning was performed with supportive rolls under shoulders and pelvis providing a free abdomen in order to minimized increases in intra-abdominal pressure.

**Study protocol**


The FIO<sub>2</sub> (1.0) remained unchanged throughout the experiment. A 20-min equilibration period was given for each modification following the study protocol. After time point T<sub>0</sub> the standard ventilator was replaced by an oscillatory ventilator (Sensormedics 3100B, Viasys, Conshohocken, PA, USA) without changes in P<sub>mean</sub>. The animals were randomly assigned to two groups (n = 7 each):

Group 1 : HFOV (Bias flow = 30 l/min, amplitude = 70 cm H<sub>2</sub>O, I:E = 1:1, RR = 300/min).

Group 2 : HFOV (Bias flow = 30 l/min, amplitude = 70 cm H<sub>2</sub>O, I:E = 1:1, RR = 300/min) and prone positioning.

A 20-min period was given for equilibration between each modification and followed by measurements of hemodynamics, blood gases and respiratory parameters. The following modifications were performed after completion of measurements terminating the previous 20-min period (Figure 1):

- (1) T<sub>baseline</sub>: 30 min after instrumentation.
- (2) T<sub>0</sub>: 60 min after last pulmonary lavage.
- (3) T<sub>20</sub>: HFOV (P<sub>mean</sub> = 16 cmH<sub>2</sub>O) had been started in both groups



**Figure 2**

Arterial oxygen partial pressure ( $\text{PaO}_2$ ) throughout the study protocol. Data are mean  $\pm$  standard deviation. #  $p < 0.05$  vs.  $T_0$ ; \*  $p < 0.05$  vs.  $T_{T-20}$ ; §  $p < 0.05$  HFOV vs. HFOV prone

(4)  $T_{40}$ : The animals had been positioned prone in group 2; no changes were made in group 1.

(5)  $T_{60}$ :  $P_{\text{mean}}$  had been increased from 16 to 19  $\text{cmH}_2\text{O}$ .  $P_{\text{mean}'}$  measured at the tip of the endotracheal tube, was the ventilatory parameter modified during the experiment from time point  $T_{40}$  onwards. To change  $P_{\text{mean}'}$  continuous distending pressure (CDP) was elevated in steps of 3  $\text{cm H}_2\text{O}$ .

(6)  $T_{80}$ :  $P_{\text{mean}} = 22 \text{ cmH}_2\text{O}$ . (7)  $T_{100}$ :  $P_{\text{mean}} = 25 \text{ cmH}_2\text{O}$ . (8)  $T_{120}$ :  $P_{\text{mean}} = 28 \text{ cmH}_2\text{O}$ . (9)  $T_{140}$ :  $P_{\text{mean}} = 31 \text{ cmH}_2\text{O}$ . (10)  $T_{160}$ :  $P_{\text{mean}} = 34 \text{ cmH}_2\text{O}$ .

(11)  $T_{180}$ :  $P_{\text{mean}}$  had been decreased to 16  $\text{cmH}_2\text{O}$  ( $\approx T_0$ ). HFOV had been discontinued and ventilation had been set to  $T_0$ -values in all groups (PCV, PEEP = 5  $\text{cm H}_2\text{O}$ ,  $V_T = 6 \text{ ml/kg}$ , RR = 40/min). The animals in group 2 had been positioned supine.

At the end of the experiment the animals were killed using an intravenous overdose of sodium thiopental and  $T_{61}$  (Intervet, Unterschleissheim, Germany).

### Statistical analysis

Values are reported as mean  $\pm$  SD. Statistical analyses were performed with Statistica for Windows, version 5.1 (StatSoft, Tulsa, OK, USA). Two-way analysis of variance (ANOVA) for repeated measurements with factors mode


and time were used for data analysis. Student-Newman-Keuls' post hoc test was used for comparison of significant ANOVA results within and between the groups. Data of the first measurement set ( $T_{\text{baseline}}$ ) were only compared with data of the second measurement set ( $T_0$ ). P values less than 0.05 were considered significant.

### Results

Detailed data regarding hemodynamics, blood gases and respiratory parameters are presented in table 1.  $\text{PaO}_2$ -, OI- and CO-changes during the experimental period are displayed in figures 2, 3, 4.

### Lung Injury

All animals passing  $T_{\text{baseline}}$  survived the study period. Acute lung injury was induced in all animals by means of repeated lung lavages ( $19 \pm 2$ ) with significant changes ( $p < 0.05$ ) in  $\text{PaO}_2$ ,  $\text{PaCO}_2$ ,  $\text{SvO}_2$ ,  $P_{\text{mean}'}$ , PIP, Qs/Qt, CVP, MPAP and CO:  $\text{PaO}_2$  decreased from  $652 \pm 37.5 \text{ mmHg}$  to  $55 \pm 7.5 \text{ mmHg}$ ,  $\text{PaCO}_2$  increased from  $41 \pm 3 \text{ mmHg}$  to  $76.5 \pm 9 \text{ mmHg}$ ,  $\text{SvO}_2$  decreased from  $83 \pm 6.5 \%$  to  $60.5 \pm 10 \%$ ,  $P_{\text{mean}}$  increased from  $11.5 \pm 1.4 \text{ cmH}_2\text{O}$  to  $16.5 \pm 1.2 \text{ cmH}_2\text{O}$ , PIP increased from  $19 \pm 3 \text{ cmH}_2\text{O}$  to  $28 \pm 2.5 \text{ cmH}_2\text{O}$ , Qs/Qt increased from  $1 \pm 0.5 \%$  to  $60.5 \pm 11 \%$ , CVP increased from  $7.8 \pm 1.3 \text{ mmHg}$  to  $10 \pm 2.9 \text{ mmHg}$ , MPAP increased from  $22.5 \pm 5.5$  to  $30.5 \pm 5 \text{ mmHg}$ , CO increased from  $3.1 \pm 0.4 \text{ l/min}$  to  $5.7 \pm 0.8 \text{ l/min}$ . No significant differences could be detected between the 2 groups for the parameters tested at time points  $T_{\text{baseline}}$  and



**Figure 3**  
 Oxygenation Index (OI) throughout the study protocol Data are mean ± standard deviation. # p < 0.05 vs. T<sub>0</sub>; \* p < 0.05 vs. T<sub>T-20</sub>; § p < 0.05 HFOV vs. HFOV prone

T<sub>0</sub>. Reduction of P<sub>mean</sub> to the T<sub>0</sub>-level (T<sub>180</sub>) combined with PCV at T<sub>0</sub>-ventilator settings resulted in immediate significant increases in CO and Qs/Qt to T<sub>0</sub>-values in both groups without significant differences between the groups. Accordingly PaO<sub>2</sub> decreased significantly to T<sub>0</sub>-levels.

**Pulmonary gas exchange**

Oxygenation improved significantly with rising P<sub>mean</sub> in both groups. At T<sub>80</sub> and T<sub>100</sub> PaO<sub>2</sub> was significantly higher in the prone positioned animals. A significantly higher PaO<sub>2</sub> compared to the preceding time point was detected at T<sub>80</sub> in the prone positioned animals and at T<sub>120</sub> in the animals positioned supine. Significant improvement of OI occurred immediately after prone positioning (T<sub>40</sub>) lasting until T<sub>100</sub>. SvO<sub>2</sub> was significantly higher from T<sub>40</sub> onwards in the HFOV-prone group if compared to T<sub>0</sub> without detectable significant differences between the groups. All animals remained hypercapnic with a PaCO<sub>2</sub> greater 70 mmHg resulting in a pH of less than 7.23 throughout the experiment in both groups.

**Respiratory parameters**

PIP increased significantly in all groups with rising P<sub>mean</sub> without differences between the groups.

**Hemodynamics**


MAP remained stable in both groups. CVP and PCWP started to rise significantly in the HFOV group from T<sub>100</sub>

and T<sub>140</sub> respectively if compared to T<sub>0</sub>. MPAP was increased significantly in both groups from T<sub>140</sub> if compared to T<sub>0</sub>. From T<sub>40</sub> to T<sub>140</sub> CO and HR were significantly lower and continuously falling in the HFOV-prone group. At T<sub>160</sub> and T<sub>180</sub> no differences between the groups could be detected regarding CO and HR. Qs/Qt was significantly lower from T<sub>40</sub> to T<sub>120</sub> in the HFOV-prone group without differences between the groups from T<sub>140</sub> onwards.

**Discussion**

We evaluated the effects of prone positioning combination of HFOV and prone positioning in an adult animal model of ARDS. The major findings of our study are: 1) HFOV and prone positioning improves oxygenation at a lower P<sub>mean</sub> than HFOV and supine positioning. 2) HFOV and prone positioning result in significant reduction of pulmonary shunt fraction and normalisation of cardiac output at a lower P<sub>mean</sub> than HFOV and supine positioning. 3) Hypercapnia was neither ameliorated by HFOV nor combination of HFOV with prone positioning.

Since therapeutic alternatives are lacking and the underlying concepts sound reasonable, multimodal therapeutic approaches are commonly used for salvage therapy in patients with ARDS [19]. Apart from subsets of patients in other HFOV trails, the combined use of HFOV and prone positioning is described in one case report and was investigated systematically in a prospective randomized study including 39 medical patients [15,20]. Papazian et al.



**Figure 4**

Cardiac output (CO) throughout the study protocol. Data are mean ± standard deviation. # p < 0.05 vs. T<sub>0</sub>; \* p < 0.05 vs. T<sub>T</sub>. 20; § p < 0.05 HFOV vs. HFOV prone

found the prone position combined with HFOV and PCV superior to HFOV and supine positioning in terms of oxygenation, but failed to demonstrate additive effects. However, the inflammatory mediators were elevated during HFOV-prone but not during HFOV-supine. The authors themselves put these results into perspective, since a control group was lacking and a time dependent natural change in the concentration of inflammatory mediators could not be excluded. It is a limitation of our study, that we did not investigate a control group ventilated in a conventional lung protective mode and positioned prone in order to detect additive effects of the two treatment modalities. Papazian et al. stressed the difficulties associated with bronchoalveolar lavages in ARDS patients in terms of patient safety and feasibility. This calls for long term experiments with large animals comparing conventional lung protective ventilation and HFOV with and without prone positioning looking not only at gas exchange and respiratory mechanics but also at histology and inflammatory mediators.

Current concepts to ameliorate the detrimental effects of VILI focus on reduction of volutrauma, barotrauma, atelectrauma and biotrauma [21]. It was shown in a small animal model, that HFOV had the same effect on oxygenation and pulmonary compliance than a conventional lung protective ventilatory approach but also

reduced the systemic inflammatory response [22-24]. However, tracheal tube size, respiratory frequency and pressure amplitude are markedly different in small animals resulting in non-comparable changes of pulmonary mechanics and oscillatory pressure transmission. Therefore, experiments in large animals should be performed before HFOV is assessed systematically in adult patients with ARDS. Aiming to simulate a life-threatening clinical scenario, we induced an acute lung injury with severe hypoxemia and hypercapnia.

It was striking, that 19 ± 2 lavages with 30 ml/kg isotonic saline were needed to reach the targeted PaO<sub>2</sub>-value, suggesting a lung protective effect of the low-tidal-volume approach during ARDS-induction even on a low PEEP-level (5 cmH<sub>2</sub>O). Intrinsic PEEP was measured during PCV by means of an endexpiratory occlusion maneuver for five seconds after every third lavage and was always less than 1 cm H<sub>2</sub>O. In two studies using sheep with a body weight of 30 kg, 4 lavages were needed to achieve a PaO<sub>2</sub> of less than 120 mmHg. These animals were ventilated with a V<sub>T</sub> of 12 ml/kg in a volume controlled mode [25,26]. Stability of the experimentally induced ARDS was proven two-fold: 1) A stabilisation period of 60 min. with unchanged ventilatory parameters was kept between the last pulmonary lavage and T<sub>0</sub>. 2) After T<sub>160'</sub> HFOV and prone positioning were discontinued, all animals were

ventilated with PCV and PEEP was reduced to 5 cmH<sub>2</sub>O [26]. This manoeuvre resulted in immediate reversal of PaO<sub>2</sub>, PaCO<sub>2</sub> and hemodynamics to T<sub>0</sub>-values.

Although the combined application of HFOV and prone positioning improved oxygenation, normalized cardiac output and significantly reduced pulmonary shunt fraction at a lower P<sub>mean</sub> than HFOV alone, hypercapnia was not influenced in our experiment. This is consistent with clinical results, since normocapnia was not achievable with HFOV alone in many adults with ARDS [8,27]. The ability to control the PaCO<sub>2</sub> with the least possible ventilator pressure amplitude, e.g. by using HFOV, might result in further lung protection [28]. However, V<sub>T</sub>-reduction increases the risk of hypercapnia, thereby aggravating the pulmonary inflammatory response [29]. Even though permissive hypercapnia does not increase mortality and might have beneficial effects, such as lung protection from reperfusion injury [30], there are clinical situations where hypercapnia is contraindicated [31]. We knew from pilot experiments that CO<sub>2</sub>-elimination could only be increased in our animal model using a respiratory rate of less than 3 Hz, losing the advantage of oscillation in reducing lung damage [32]. Another possibility to improve CO<sub>2</sub>-elimination during HFOV is deflation of the cuff of the endotracheal tube. Since comparability of P<sub>mean</sub> was a prerequisite for the study and airway-pressure measurement at the tip of the endotracheal tube are prone to artefacts, we could not realize this option. There might be a need to combine HFOV with extracorporeal CO<sub>2</sub>-elimination whenever normocapnia is mandatory, e.g. in patients with cerebral oedema [33].

A lung volume between the lower and upper inflection point, derived from the inflation pressure-volume curve, is traditionally interpreted as ideal for oxygenation [34]. Evidence is raising, that best oxygenation is a better indicator for an open lung-PEEP during a decremental PEEP-trial after a recruitment manoeuvre [35]. We aimed to achieve optimal lung volume using increases in PaO<sub>2</sub> as a crude surrogate for alveolar recruitment and avoidance of hyperinflation. Since an initial recruitment manoeuvre in our animals might have resulted in fatal cardiovascular collapse and barotrauma [23], we increased P<sub>mean</sub> stepwise and did not create a pressure volume relationship [36,37]. Generation of a pressure volume curve can cause lung derecruitment [38]. Further deterioration of oxygenation in our animals would have implied a high risk of irreversible hypoxia. The severity of hypoxemia present in our animals after ARDS-induction was our motivation not to randomize the airway pressure changes but to stepwise increase P<sub>mean</sub>[39].

It is the major limitation of this study, that it was performed in pigs and not in patients. In adults not surfactant

deficiency but alveolar flooding is the predominant mechanism in ARDS-development. It limits the transferability of most ARDS model derived results to clinical practice [40].

## Conclusion

In this saline lavage induced porcine model of ARDS, we showed in a clinically relevant scenario, that the combination of HFOV and prone positioning improved oxygenation at a lower P<sub>mean</sub> than HFOV combined with supine positioning. In addition, reduction of the pulmonary shunt fraction and normalisation of the cardiac output was achieved at lower airway pressures. The ventilator pressure amplitude is a major determinant of VILI. HFOV might be a step towards further lung protection, since sufficient oxygenation can be restored or maintained with a significant reduction of the ventilator pressure amplitude when compared to standard respirator modes. However, HFOV failed to be a major component in ARDS treatment algorithms in adult patients. Having in view a long history of failed multimodal treatment approaches in ARDS research, we now conclude from our results that a combination of HFOV and prone positioning seems promising and should be further investigated systematically and compared to conventional lung protective ventilation. Long term trials in large animals and acquisition of histologic and immunologic data clearly seem justified.

## Abbreviations

- ARDS acute respiratory distress syndrome
- CaO<sub>2</sub> arterial oxygen content
- CCO<sub>2</sub> pulmonary capillary oxygen content
- CDP continuous distending pressure
- CO cardiac output
- CvO<sub>2</sub> mixed venous oxygen content
- CVP central venous pressure
- FIO<sub>2</sub> fraction of inspired oxygen
- HFOV high frequency oscillatory ventilation
- HR heart rate
- I:E inspiratory to expiratory ratio
- MAP mean arterial pressure
- MPAP mean pulmonary artery pressure

OI Oxygenation index

PaCO<sub>2</sub> Arterial carbon dioxide partial pressure

PaO<sub>2</sub> Arterial oxygen partial pressure

PEEP positive end-expiratory pressure

PIP peak inspiratory pressure

PCV pressure controlled ventilation

PCWP pulmonary capillary wedge pressure

P<sub>mean</sub> mean airway pressure

Qs/Qt pulmonary shunt fraction

RR; f respiratory rate

V<sub>T</sub> tidal volume

### Competing interests

The author(s) declare that they have no competing interests.

### Authors' contributions

JB designed and coordinated the study and drafted the manuscript. JB and RM performed the experiments. MK performed the statistical analysis. MA, CG and NR participated in the design and coordination of the study and helped to draft the manuscript. All authors read and approved the final manuscript.

### Acknowledgements

The authors thank Alois Reichert (Dept. of Surgery) for technical assistance, Brigitte Paul and Juergen Schoell (Dept. of Anesthesiology and Intensive Care Medicine) for logistic and Karin Ulrichs, Ph.D. (Dept. of Surgery) for organisatory support.

The study was conducted with departmental funding. The high frequency oscillatory ventilator was provided by Viasys Healthcare, Hoechberg, Wuerzburg, Germany.

### References

1. **Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The Acute Respiratory Distress Syndrome Network.** *N Engl J Med* 2000, **342**:1301-1308.
2. Fan E, Needham DM, Stewart TE: **Ventilatory management of acute lung injury and acute respiratory distress syndrome.** *JAMA* 2005, **294**:2889-2896.
3. Imai Y, Parodo J, Kajikawa O, de Perrot M, Fischer S, Edwards V, Cutz E, Liu M, Keshavjee S, Martin TR, Marshall JC, Ranieri VM, Slutsky AS: **Injurious mechanical ventilation and end-organ epithelial cell apoptosis and organ dysfunction in an experimental model of acute respiratory distress syndrome.** *JAMA* 2003, **289**:2104-2112.
4. Luhr OR, Karlsson M, Thorsteinsson A, Rylander C, Frostell CG: **The impact of respiratory variables on mortality in non-ARDS and ARDS patients requiring mechanical ventilation.** *Intensive Care Medicine* 2000, **26**:508-517.
5. Bone RC, Maunder R, Slotman G, Silverman H, Hyers TM, Kerstein MD, Ursprung JJ: **An early test of survival in patients with the adult respiratory distress syndrome. The PaO<sub>2</sub>/Fio<sub>2</sub> ratio and its differential response to conventional therapy. Prostaglandin E1 Study Group.** *Chest* 1989, **96**:849-851.
6. Hemmila MR, Rowe SA, Boules TN, Miskulin J, McGillicuddy JW, Schuerer DJ, Haft JW, Swaniker F, Arbabi S, Hirschi RB, Bartlett RH: **Extracorporeal life support for severe acute respiratory distress syndrome in adults.** *Ann Surg* 2004, **240**:595-605.
7. Derdak S: **High-frequency oscillatory ventilation for acute respiratory distress syndrome in adult patients.** *Crit Care Med* 2003, **31**:S317-S323.
8. Derdak S, Mehta S, Stewart TE, Smith T, Rogers M, Buchman TG, Carlin B, Lowson S, Granton J: **High-frequency oscillatory ventilation for acute respiratory distress syndrome in adults: a randomized, controlled trial.** *Am J Respir Crit Care Med* 2002, **166**:801-808.
9. Gattinoni L, Tognoni G, Pesenti A, Taccone P, Mascheroni D, Labarta V, Malacrida R, Di Giulio P, Fumagalli R, Pelosi P, Brazzi L, Latini R: **Effect of prone positioning on the survival of patients with acute respiratory failure.** *N Engl J Med* 2001, **345**:568-573.
10. Vieillard-Baron A, Rabiller A, Chergui K, Peyrouset O, Page B, Beauchet A, Jardin F: **Prone position improves mechanics and alveolar ventilation in acute respiratory distress syndrome.** *Intensive Care Med* 2005, **31**:220-226.
11. Gattinoni L, Vagginelli F, Carlesso E, Taccone P, Conte V, Chiumello D, Valenza F, Caironi P, Pesenti A: **Decrease in PaCO<sub>2</sub> with prone position is predictive of improved outcome in acute respiratory distress syndrome.** *Crit Care Med* 2003, **31**:2727-2733.
12. Gannier M, Michelet P, Thirion X, Arnal JM, Sainty JM, Papazian L: **Prone position and positive end-expiratory pressure in acute respiratory distress syndrome.** *Crit Care Med* 2003, **31**:2719-2726.
13. Pelosi P, Brazzi L, Gattinoni L: **Prone position in acute respiratory distress syndrome.** *Eur Respir J* 2002, **20**:1017-1028.
14. Varkul MD, Stewart TE, Lapinsky SE, Ferguson ND, Mehta S: **Successful use of combined high-frequency oscillatory ventilation, inhaled nitric oxide, and prone positioning in the acute respiratory distress syndrome.** *Anesthesiology* 2001, **95**:797-799.
15. Papazian L, Gannier M, Marin V, Donati S, Arnal JM, Demory D, Roch A, Forel JM, Bongrand P, Bregeon F, Sainty JM: **Comparison of prone positioning and high-frequency oscillatory ventilation in patients with acute respiratory distress syndrome.** *Crit Care Med* 2005, **33**:2162-2171.
16. Nagano O, Fujii H, Morimatsu H, Mizobuchi S, Goto K, Katayama H, Hirakawa M, Yamada Y: **An adult with ARDS managed with high-frequency oscillatory ventilation and prone position.** *J Anesth* 2002, **16**:75-78.
17. Hallman M, Merritt TA, Jarvenpaa AL, Boynton B, Mannino F, Gluck L, Moore T, Edwards D: **Exogenous human surfactant for treatment of severe respiratory distress syndrome: a randomized prospective clinical trial.** *J Pediatr* 1985, **106**:963-969.
18. Stenqvist O: **Practical assessment of respiratory mechanics.** *Br J Anaesth* 2003, **91**:92-105.
19. Fan E, Mehta S: **High-frequency oscillatory ventilation and adjunctive therapies: inhaled nitric oxide and prone positioning.** *Crit Care Med* 2005, **33**:S182-S187.
20. Krishnan JA, Brower RG: **High-frequency ventilation for acute lung injury and ARDS.** *Chest* 2000, **118**:795-807.
21. Imai Y, Slutsky AS: **High-frequency oscillatory ventilation and ventilator-induced lung injury.** *Crit Care Med* 2005, **33**:S129-S134.
22. Imai Y, Kawano T, Miyasaka K, Takata M, Imai T, Okuyama K: **Inflammatory Chemical Mediators During Conventional Ventilation and During High-Frequency Oscillatory Ventilation.** *American Journal of Respiratory and Critical Care Medicine* 1994, **150**:1550-1554.
23. Imai Y, Nakagawa S, Ito Y, Kawano T, Slutsky AS, Miyasaka K: **Comparison of lung protection strategies using conventional and high-frequency oscillatory ventilation.** *Journal of Applied Physiology* 2001, **91**:1836-1844.
24. Kerr CL, Veldhuizen RAW, Lewis JF: **Effects of high-frequency oscillation on endogenous surfactant in an acute lung injury**

- model.** *American Journal of Respiratory and Critical Care Medicine* 2001, **164**:237-242.
25. Goddon S, Fujino Y, Hromi JM, Kacmarek RM: **Optimal mean airway pressure during high-frequency oscillation - Predicted by the pressure-volume curve.** *Anesthesiology* 2001, **94**:862-869.
  26. Sedek KA, Takeuchi M, Suchodolski K, Vargas SO, Shimaoka M, Schnitzer JJ, Kacmarek RM: **Open-lung protective ventilation with pressure control ventilation, high-frequency oscillation, and intratracheal pulmonary ventilation results in similar gas exchange, hemodynamics, and lung mechanics.** *Anesthesiology* 2003, **99**:1102-1111.
  27. David M, Weiler N, Heinrichs W, Neumann M, Joost T, Markstaller K, Eberle B: **High-frequency oscillatory ventilation in adult acute respiratory distress syndrome.** *Intensive Care Med* 2003, **29**:1656-1665.
  28. Frank JA, Gutierrez JA, Jones KD, Allen L, Dobbs L, Matthay MA: **Low tidal volume reduces epithelial and endothelial injury in acid-injured rat lungs.** *Am J Respir Crit Care Med* 2002, **165**:242-249.
  29. Lang JD, Figueroa M, Sanders KD, Aslan M, Liu Y, Chumley P, Freeman BA: **Hypercapnia via reduced rate and tidal volume contributes to lipopolysaccharide-induced lung injury.** *Am J Respir Crit Care Med* 2005, **171**:147-157.
  30. Laffey JG, O'Croinin D, McLoughlin P, Kavanagh BP: **Permissive hypercapnia--role in protective lung ventilatory strategies.** *Intensive Care Med* 2004, **30**:347-356.
  31. Doerr CH, Gajic O, Berrios JC, Caples S, Abdel M, Lymp JF, Hubmayr RD: **Hypercapnic Acidosis Impairs Plasma Membrane Wound Resealing in Ventilator-injured Lungs.** *Am J Respir Crit Care Med* 2005, **171**:1371-1377.
  32. Sedek KA, Takeuchi M, Suchodolski K, Kacmarek RM: **Determinants of tidal volume during high-frequency oscillation.** *Crit Care Med* 2003, **31**:227-231.
  33. David M, Heinrichs W: **High-frequency oscillatory ventilation and an interventional lung assist device to treat hypoxaemia and hypercapnia.** *Br J Anaesth* 2004, **93**:582-586.
  34. McCulloch PR, Forkert PG, Froese AB: **Lung volume maintenance prevents lung injury during high frequency oscillatory ventilation in surfactant-deficient rabbits.** *Am Rev Respir Dis* 1988, **137**:1185-1192.
  35. Hickling KG: **Best compliance during a decremental, but not incremental, positive end-expiratory pressure trial is related to open-lung positive end-expiratory pressure: a mathematical model of acute respiratory distress syndrome lungs.** *Am J Respir Crit Care Med* 2001, **163**:69-78.
  36. Luecke T, Meinhardt JP, Herrmann P, Weisser G, Pelosi P, Quintel M: **Setting mean airway pressure during high-frequency oscillatory ventilation according to the static pressure-volume curve in surfactant-deficient lung injury: a computed tomography study.** *Anesthesiology* 2003, **99**:1313-1322.
  37. Rimensberger PC, Cox PN, Frndova H, Bryan AC: **The open lung during small tidal volume ventilation: concepts of recruitment and "optimal" positive end-expiratory pressure.** *Crit Care Med* 1999, **27**:1946-1952.
  38. Henzler D, Mahnken A, Dembinski R, Waskowiak B, Rossaint R, Kuhlén R: **Repeated generation of the pulmonary pressure-volume curve may lead to derecruitment in experimental lung injury.** *Intensive Care Med* 2005, **31**:302-310.
  39. Krishnan RK, Meyers PA, Worwa C, Goertz R, Schauer G, Mammel MC: **Standardized lung recruitment during high frequency and conventional ventilation: similar pathophysiologic and inflammatory responses in an animal model of respiratory distress syndrome.** *Intensive Care Med* 2004, **30**:1195-1203.
  40. Kloot TE, Blanch L, Melyne YA, Weinert C, Adams AB, Marini JJ, Shapiro RS, Nahum A: **Recruitment maneuvers in three experimental models of acute lung injury. Effect on lung volume and gas exchange.** *Am J Respir Crit Care Med* 2000, **161**:1485-1494.

### Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2253/6/4/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

