

CORRECTION

Open Access



Correction to: Incidence and severity of postoperative sore throat: a randomized comparison of Glidescope with Macintosh laryngoscope

Mansoor Aqil^{1*}, Mueen Ullah Khan¹, Saara Mansoor², Saad Mansoor², Rashid Saeed Khokhar¹ and Abdul Sattar Narejo³

Correction

We are thankful to Dr. Deepak Gupta (Clinical Assistant Professor Anesthesiology, Wayne State University/Detroit Medical Center) for bringing to our attention a typographical error in our manuscript [1].

Table 2, ROW 12 was wrongly written as:

Number of patients with sore throat (Yes/No)

At 0 h 22/48 (Glidescope), 41/48 (Macintosh), 0.001 (*P* value).

It should be read as: Number of patients with sore throat (Yes/No)

At 0 h 22/48 (Glidescope), 41/29 (Macintosh), 0.001 (*P* value).

Author details

¹Department of Anesthesiology, King Saud University Medical City, P.O Box 7805, Riyadh 11472, Saudi Arabia. ²College of Medicine, Alfaisal University, Riyadh, Saudi Arabia. ³Department of Anesthesiology, King Khalid Hospital, Hail, Saudi Arabia.

Published online: 17 August 2018

Reference

1. Aqil M, Khan MU, Mansoor S, Mansoor S, Khokhar RS, Narejo AS. Incidence and severity of postoperative sore throat: a randomized comparison of Glidescope with Macintosh laryngoscope. *BMC Anesthesiol.* 2017;17:127. <https://doi.org/10.1186/s12871-017-0421-4>.

* Correspondence: mansooraqil@yahoo.com

¹Department of Anesthesiology, King Saud University Medical City, P.O Box 7805, Riyadh 11472, Saudi Arabia

