

CORRECTION

Open Access



Correction to: Postoperative delirium in critically ill surgical patients: incidence, risk factors, and predictive scores

Onuma Chaiwat^{1,3*}, Mellada Chanidnuan¹, Worapat Pancharoen¹, Kittiya Vijitmal², Praniti Danpornprasert², Puriwat Toaditthep¹ and Chayanan Thanakiattiwibun³

Correction to: *BMC Anesthesiology* (2019) 19:39

<https://doi.org/10.1186/s12871-019-0694-x>

Following publication of the original article [1], the authors reported a missing data on Table 1 in their paper. The original article [1] has been updated.

The correct Table 1 is shown below.

Author details

¹Department of Anesthesiology, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand. ²Division of Critical Care Medicine, Department of Medicine, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok, Thailand. ³Integrated Perioperative Geriatric Excellent Research Center, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok, Thailand.

Published online: 22 April 2019

Reference

1. Chaiwat, et al. Postoperative delirium in critically ill surgical patients: incidence, risk factors, and predictive scores. *BMC Anesthesiology*. 2019;19:39. <https://doi.org/10.1186/s12871-019-0694-x>.

* Correspondence: onuma.cha@mahidol.ac.th

¹Department of Anesthesiology, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand

³Integrated Perioperative Geriatric Excellent Research Center, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok, Thailand

Full list of author information is available at the end of the article



Table 1 Baseline characteristics of delirious and non-delirious patients

Variables	Delirium (n = 61)	No delirium (n = 189)	P-value
Demographic data			
Age (years)	72.7 ± 11.4	61.4 ± 16.8	< 0.001
≥ 60	55 (90.2%)	114 (60.3%)	< 0.001
Gender			
Male	31 (50.8%)	90 (47.6%)	0.768
Comorbidities			
Hypertension	50 (81.9%)	105 (55.6%)	< 0.001
DM	26 (42.6%)	37 (19.6%)	0.001
Cardiac disease	21 (34.4%)	43 (22.8%)	0.091
Previous stroke	15 (24.6%)	18 (9.5%)	0.004
Modified IQCODE score ≥ 3.42	10 (16.39%)	6 (3.2%)	0.001
ESRD or CKD stage 4–5	10 (16.4%)	24 (12.7%)	0.520
Cirrhosis	3 (4.9%)	9 (4.8%)	1.000
Smoking history pack year	41.9 ± 27.1	24.4 ± 21.6	0.155
≥ 30 pack year	10 (16.4%)	20 (10.6%)	0.259
Current alcohol consumption	6 (9.8%)	11 (5.8%)	0.378
Coma	17 (27.9%)	16 (8.5%)	< 0.001
Intraoperative data			
Emergency Surgery	34 (55.74%)	74 (39.2%)	0.026
Vascular surgery	20 (32.8%)	32 (16.9%)	0.011
Non-vascular surgery			
Intra-abdominal	23 (37.7%)	65 (34.4%)	0.646
Orthopedic	3 (4.9%)	26 (13.8%)	0.068
Gynecological	3 (4.9%)	23 (12.2%)	0.147
Other	12 (19.7%)	43 (22.8%)	0.723
Operation time	193.5 ± 162.6	234.8 ± 178.9	0.111
Intraoperative blood loss (mL)	250 (60–700)	400 (100–1400)	0.079
Intraoperative PRC transfusion (mL)	264 (0–663)	0 (0–1023)	0.865
Hypoxia	3 (4.9%)	7 (3.7%)	0.710
Intraoperative hypotension	50 (82.0%)	146 (77.3%)	0.480
ICU admission			
Hemoglobin (mg/dL)	10.5 ± 2.5	10.9 ± 2.1	0.156
Blood sugar at admission (mg/dL)	155.6 ± 42.1	152.2 ± 46.3	0.611
Serum albumin (mg/dL)	2.7 ± 0.6	2.8 ± 0.7	0.168
BUN/Cr ratio > 20	22 (36.1%)	48 (25.4%)	0.139
Sepsis	24 (39.3%)	37 (19.6%)	0.003
APACHE II score	12.1 ± 4.8	8.3 ± 3.9	< 0.001
SOFA score	5.8 ± 3.4	3.4 ± 2.7	< 0.001
Mechanical ventilation	54 (88.5%)	131 (69.2%)	0.002
Medication			
Preoperative-Benzodiazepine	8 (13.1%)	10 (5.3%)	0.049

Table 1 Baseline characteristics of delirious and non-delirious patients (Continued)

Variables	Delirium (n = 61)	No delirium (n = 189)	P-value
Intraoperative-Benzodiazepine	32 (53.3%)	93 (47.7%)	0.658
Benzodiazepine (Pre and Postoperative)	27 (44.3%)	36 (19.1%)	< 0.001
Preoperative-Opioid	5 (8.2%)	18 (9.5%)	1.000
Intraoperative-Opioid	58 (96.7%)	176 (94.1%)	0.740
Opioid (Pre and Postoperative)	59 (96.7%)	184 (97.4%)	1.000
Postoperative-Propofol use	20 (32.8)	31 (16.4%)	0.010

ESRD End stage renal disease, CKD Chronic kidney disease, DM Diabetes mellitus, Modified IQCODE score Modified informant questionnaire on cognitive decline in the elderly score, BUN Blood urea nitrogen, Cr Creatinine, APACHE II score Acute physiology and chronic health evaluation II score, SOFA score Sequential organ failure assessment score
Data presented as mean ± SD or median (IQR) or N (%)