

CORRECTION

Open Access



Correction to: Multimodal analgesia with ropivacaine wound infiltration and intravenous flurbiprofen axetil provides enhanced analgesic effects after radical thyroidectomy: a randomized controlled trial

Xiaoxi Li, Ling Yu, Jiaonan Yang and Hongyu Tan*

Correction to: BMC Anesthesiol (2019) 19:167
<https://doi.org/10.1186/s12871-019-0835-2>

Following publication of the original article [1], the authors reported that there has been a mistake regarding the unit of duration of surgery, consciousness recovery time, and extubation time in Table 1; the correct unit is minute. This is the correct information in Table 1:

Duration of surgery (min)
Consciousness recovery time (min)
Extubation time (min)

Published online: 13 November 2019

Reference

1. Li X, et al. Multimodal analgesia with ropivacaine wound infiltration and intravenous flurbiprofen axetil provides enhanced analgesic effects after radical thyroidectomy: a randomized controlled trial. *BMC Anesthesiol*. 2019; 19:167. <https://doi.org/10.1186/s12871-019-0835-2>.

* Correspondence: maggitan@163.com

Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Department of Anesthesiology, Peking University Cancer Hospital & Institute, #52 Fucheng Street, Haidian District, Beijing 100142, China

