CORRECTION Open Access

## Correction to: Effects of different sufentanil target concentrations on the MAC<sub>BAR</sub> of sevoflurane in patients with carbon dioxide pneumoperitoneum stimulus



Yanxia Guo<sup>1†</sup>, Dan Wang<sup>1†</sup>, Xiaolin Yang<sup>1\*</sup>, Pingping Jiang<sup>1</sup>, Juan Xu<sup>1</sup> and Guoyuan Zhang<sup>2</sup>

Correction to: BMC Anesthesiology 20, 239 (2020) https://doi.org/10.1186/s12871-020-01160-1

Following publication of the original article [1], the authors reported an error in Fig. 1. The correct version of Fig. 1 is shown below.

## Author details

<sup>1</sup>Department of Anaesthesiology, Affiliated Hospital of North Sichuan Medical College, Nanchong 637000, Sichuan, China. <sup>2</sup>Department of Clinical Laboratory, Affiliated Hospital of North Sichuan Medical College, Nanchong 637000, Sichuan, China.

Published online: 23 October 2020

## Reference

 Guo Y, Wang D, Yang X, et al. Effects of different sufentanil target concentrations on the MAC<sub>BAR</sub> of sevoflurane in patients with carbon dioxide pneumoperitoneum stimulus. BMC Anesthesiol. 2020;20:239. https:// doi.org/10.1186/s12871-020-01160-1.

The original article can be found online at https://doi.org/10.1186/s12871-020-01160-1.

<sup>&</sup>lt;sup>1</sup>Department of Anaesthesiology, Affiliated Hospital of North Sichuan Medical College, Nanchong 637000, Sichuan, China




© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

<sup>\*</sup> Correspondence: 879921874@qq.com

<sup>&</sup>lt;sup>†</sup>Yanxia Guo and Dan Wang contributed equally to this article and share first authorship

Guo et al. BMC Anesthesiology (2020) 20:269 Page 2 of 2



**Fig. 1** Dixon up-and-down plots for each group. The plasma target concentration of sufentanil in groups  $S_0$ ,  $S_1$ ,  $S_2$ ,  $S_3$  and  $S_4$  was 0.0, 0.1, 0.3, 0.5 and 0.7 ng ml<sup>-1</sup>, respectively. The empty (solid) circle represents the negative (positive) reaction to hemodynamics parameters, and the triangle indicates the intersection of negative and positive reactions. The ninth patient was given the same concentration of sevoflurane both in group  $S_2$  and group  $S_3$ . To get six crossovers, 14, 14, 18, 20 and 15 patients were needed in groups  $S_0$ - $S_4$ , respectively